## Breast Reconstruction using tissue from the thigh (TMG-FLAP)

The tissue on the inside of the upper thigh is the second most common procedure used for breast reconstruction and is also suitable for reconstruction of both breasts. This technique is used when there is insufficient tissue on the abdomen, or the abdominal blood vessels cannot be used due to previous surgery.

The tissue from the upper inner thigh is transferred to the breast, and the blood vessels of the tissue are connected to the chest using a microscope (microsurgery).

QUESTIONS YOU MAY WANT TO ASK DURING THE CONSULTATION:

How does my breast look and feel afterwards?

When can I return to my normal daily routine?

How long will I be restricted in sport and exercise after

surgery?

Will the oncological follow-up be affected by this

technique of reconstruction?

## INFORMATION ABOUT THE PROCEDURE:


Duration of case: 3.5 hours

General Anaesthesia required

Post-operative recovery approximately 3-4 weeks

No physical activity/sport for approximately 4-6 weeks

Correspondence Address: The London Clinic, Consulting Rooms 5 Devonshire Place London W1G 6HL Private Consultancy: The London Clinic Consulting Rooms 5 Devonshire Place London W1G 6HL Contact details: +44 754 054 9873 www.farhadi.com Email: farhadij@me.com



## Prof Farhadi